

Integration formulas for Brownian motion on classical groups

Antoine Dahlqvist *

December 21, 2012

Abstract

We give new formulas for the moments of the entries of a random matrix whose law is a Brownian motion on a classical compact group. As we let the time go to infinity, these formulas imply the one B. Collins and P. Śniady obtained for Haar measure in [2].

1 Introduction

Let G be a compact group that belongs to one of the classical series $U(N)$, $O(N)$ and $USp(N)$. For any of these groups, let us denote by V their fundamental representation, that is $\mathbb{C}^N$ for $U(N)$ and $O(N)$, and $\mathbb{C}^{2N}$ for $USp(N)$. In the following, we are interested in finding an explicit expression for

$$\int_G f(g) \mu(dg) \quad (1)$$

where f is the composition of a polynomial function on $\text{End}(V)$ with the fundamental representation and μ is the measure associated to the law of a Brownian motion that we shall define in Section 3.2. Such quantities have been studied when μ is the Haar measure by B. Collins in [3] and later with P. Śniady in [2], see also [1] for a recent point of view. The subject of this article is to give formulas for (1) that are deformations of the ones obtained in [2]. Besides, our work completes an article of T. Lévy [7] who computes such integrals but for a smaller class of polynomials. One of the motivations of our computation is the fact that the law of the Brownian motion is invariant by conjugation so that integrating with respect to it defines invariant quantities for the group G . For any space E on which G acts, we will denote by E^G the set of points in E fixed by the action of G . If W is a representation of G , we denote by $\text{End}_G(W)$ the space of endomorphisms commuting with G . We will consider here the representations $W = V^{\otimes n} \otimes V^{*\otimes m}$. For such representations, $\text{End}_G(W)$ is described by the Schur-Weyl duality (see [6]). As an example, for $G = U(N)$, $\text{End}_G(V^{\otimes n})$ is the image of $\mathbb{C}[\mathfrak{S}_n]$ by the classical representation on $V^{\otimes n}$ given by permutation of the tensors.

One nice feature of our formulas is the formulation in terms of a commutative family of elements of $\mathbb{C}[\mathfrak{S}_n]$ called the Jucy-Murphy elements, whose joint spectrum is explicitly known (see [9]). For any non-zero integer i , we let $X_i = (1 \ i) + (2 \ i) + \dots + (i-1 \ i)$ be i -th Jucy-Murphy

*Laboratoire de Probabilités et Modèles Aléatoires, 4, Place Jussieu, 75252 PARIS Cedex 05. *E-mail adress:* antoine.dahlqvist@upmc.fr

element. We set $X_0 = 0$. We get a generalization for the Brownian motion of the formulas of [10].

In the first section of this article, we give a proof of an integration formula for the Haar measure on the symplectic group and recall the formulas for the orthogonal and the unitary groups. In the second Section, we reformulate this expression in the algebras $\text{End}_G(V^{\otimes n})$ and state the formulas we want to prove for the Brownian motion in Theorem 3.9 and show they are indeed a deformation of the formulas for the Haar measure. In the last Section, we give a proof of Theorem 3.9.

2 Integration formulas for the Haar measure

We give here a proof of the integration formula with respect to Haar measure on the symplectic group and recall the formulas for $U(N)$ and $O(N)$ (a proof for the compact symplectic group is also given in [4]).

Let us introduce ω the skew-symmetric bilinear form on V whose matrix in the canonical basis is $J = \begin{pmatrix} 0 & I_N \\ -I_N & 0 \end{pmatrix}$. Let us denote by $\text{USp}(2N) = \{S \in U(2N) : {}^t S J S = J\}$ the group of unitary symplectic matrices. We shall write dO, dU and dS for the Haar measure on the groups $U(N), O(N)$ and $\text{USp}(N)$. In the following we shall compute the following integral:

$$\int_{\text{USp}(2N)} S_{i_1, j_1} S_{i_2, j_2} \cdots S_{i_n, j_n} dS. \quad (2)$$

Observe that for any $S \in \text{USp}(2N)$, $\bar{S} = -JSJ$, therefore we can deduce from our computation formulas for integrals of polynomials in entries of S and $\bar{S}$. Since the Haar measure is invariant by multiplication by $-\text{Id}$, this integral is zero for odd n . Let us introduce p the endomorphism of $V^{\otimes 2n}$ defined by $p = \int_{\text{USp}(2N)} S^{\otimes 2n} dS$. Computing (2) amounts to computing

$$\langle e_{i_1} \otimes \cdots \otimes e_{i_n}, p(e_{j_1} \otimes \cdots \otimes e_{j_n}) \rangle.$$

Using invariance of the Haar measure, it is easily seen that p is an orthogonal projection onto the vector space $(V^{\otimes 2n})^{\text{USp}(2N)}$. Let $(w_l)_{l \in L}$ be any generating family of $(V^{\otimes 2n})^{\text{Sp}(2N)}$, $G = (\langle w_i, w_j \rangle)_{i, j \in L}$ its Gram matrix and $\tilde{G}$ a matrix satisfying $G\tilde{G}G = G$. The projection onto $(V^{\otimes 2n})^{\text{USp}(2N)}$ can be reformulated as

$$p = \sum_{i, j \in L} \tilde{G}_{i, j} \langle w_j, \cdot \rangle w_i. \quad (3)$$

We shall now give a generating family of $(V^{\otimes 2n})^{\text{USp}(2N)}$. Let $\mathcal{M}(2n)$ be the set of partitions by pairs of $\{1, \dots, 2n\}$. Let us denote by $\pi_0 \in \mathcal{M}(2n)$ the partition $\{\{1, 2\}, \{3, 4\}, \dots, \{2n-1, 2n\}\}$ and

$$w_{\pi_0} = \sum_{1 \leq i_1, i_2, \dots, i_{2n} \leq N} \left(\prod_{k=1}^n J_{i_{2k-1}, i_{2k}} \right) e_{i_1} \otimes e_{i_2} \otimes \cdots \otimes e_{i_{2n-1}} \otimes e_{i_{2n}}.$$

The element w_{π_0} is fixed by the action of $\text{USp}(2N)$. Let us denote by $\varepsilon : \mathfrak{S}_{2n} \rightarrow \{-1, 1\}$ the signature homomorphism and $\rho_\varepsilon : \mathfrak{S}_{2n} \rightarrow \text{GL}(V^{\otimes 2n})$ the homomorphism defined by setting for any $\sigma \in \mathfrak{S}_{2n}$ and $v_1, \dots, v_{2n} \in V$,

$$\rho_\varepsilon(\sigma) \cdot (v_1 \otimes \cdots \otimes v_{2n}) = \varepsilon(\sigma) v_{\sigma^{-1}(1)} \otimes \cdots \otimes v_{\sigma^{-1}(2n)}.$$

Let us denote by $\mathcal{H}_n$ the stabilizer of π_0 under the natural action of $\mathfrak{S}_{2n}$ on $\mathcal{M}(2n)$. The set $\mathcal{M}(2n)$ is in bijection with $\mathfrak{S}_{2n}/\mathcal{H}_n$. The group $\mathcal{H}_n$ stabilizes the element w_{π_0} under the action of $\mathfrak{S}_{2n}$ through ρ_ε , therefore for any $\pi \in \mathcal{M}(2n)$ and $\sigma \in \mathfrak{S}_{2n}$ such that $\sigma\pi_0 = \pi$, the following vector is well defined

$$w_\pi = \rho_\varepsilon(\sigma).w_{\pi_0}.$$

The action of $U(2N)$ on $V^{\otimes 2n}$ commutes with that of $\mathfrak{S}_{2n}$, therefore for any $\pi \in \mathcal{M}_{2n}$, $w_\pi \in (V^{\otimes 2n})^{\text{Sp}(2N)}$. The first geometric Theorem of invariant theory claims that all invariants are linear combinations of this ones.

Theorem 2.1 ([6] Thm. 5.2.2). *The family $(w_\pi)_{\pi \in \mathcal{M}(2n)}$ generates $(V^{\otimes 2n})^{\text{USp}(2N)}$.*

We shall compute the Gram matrix of this family. For any pair of partitions π, η of $\{1, \dots, 2n\}$, Let us denote by $\pi \vee \eta$ the finest partition coarser than π and η . Let $\#\pi$ the number of blocks of π . For any permutation $\sigma \in \mathfrak{S}_{2n}$ and any $I = (i_1, \dots, i_{2n})$ list of $2n$ integers, let us write

$$\text{inv}_2(\sigma, I) = \#\{k \in \{1, \dots, n\} : i_{\sigma(2k)} < i_{\sigma(2k-1)}\}.$$

Lemma 2.2. *i) For any $\pi, \eta \in \mathcal{M}(2n)$,*

$$\langle w_\pi, w_\eta \rangle = (-1)^n (-2N)^{\#\pi \vee \eta}.$$

ii) Let π be an element of $\mathcal{M}(2n)$, and $1 \leq i_1, \dots, i_{2n} \leq N$. If $i_l = i_k + N \pmod{2N}$, as soon as $l \sim_\pi k$, then

$$\langle w_\pi, e_{i_1} \otimes \dots \otimes e_{i_{2n}} \rangle = \varepsilon(\sigma_\pi) (-1)^{\text{inv}_2(\sigma_\pi, I)},$$

otherwise this term vanishes.

Proof. Let us prove the first point. For any partition $\{A, B\}$ of $\{1, \dots, 2n\}$ into two sets, let us say that $\{A, B\}$ forms a bipartition of $\{1, \dots, 2n\}$ compatible with a partition $\mu \in \mathcal{M}(2n)$ if each block of μ is made of an element of A and an element of B . Let π and η two elements of $\mathcal{M}(2n)$. There exists a bipartition compatible with both π and η , let us fix one $\{A, B\}$. There exists a unique $\sigma \in \mathfrak{S}_{2n}$ such that $\pi = \sigma(\eta)$ and $\sigma|_A = \text{Id}_A$. Indeed, if we identify $\mathcal{M}(2n)$ with the set of involution without fixed points of $\{1, \dots, 2n\}$, $\sigma|_B = \eta\pi$ (see figure 1). It follows that $\sigma|_B$ has exactly $\#(\pi \vee \eta)$ cycles and $\varepsilon(\sigma) = (-1)^{n - \#\pi \vee \eta}$. Let us fix a second bipartition $\{C, D\}$ compatible with π , such that

$$w_\pi = \sum_{1 \leq i_1, \dots, i_{2n}} \prod_{c \in C, d \sim_\pi c} J_{i_c, i_d} e_{i_1} \otimes \dots \otimes e_{i_{2n}}. \quad (*)$$

Let us point out that this bipartition could be non-compatible with η . On the other hand, its image $\{\sigma(C), \sigma(D)\}$ is compatible with η and

$$w_\eta = \varepsilon(\sigma) \sigma.w_\pi = \sum_{1 \leq i_1, \dots, i_{2n} \leq 2N} \varepsilon(\sigma) \prod_{c \in C, d \sim_\pi c} J_{i_{\sigma(c)}, i_{\sigma(d)}} e_{i_1} \otimes \dots \otimes e_{i_{2n}}. \quad (**)$$

Taking the scalar product of (*) and (**) gives

$$\begin{aligned} \varepsilon(\sigma) \langle w_\pi, w_\eta \rangle &= \sum_{1 \leq i_1, \dots, i_{2n} \leq 2N} \prod_{c \in C, d \sim_\pi c} J_{i_c, i_d} J_{i_{\sigma(c)}, i_{\sigma(d)}} \\ &= \sum_{1 \leq i_1, \dots, i_{2n} \leq 2N} \prod_{c \in C \cap A, d \sim_\pi c} J_{i_c, i_d} J_{i_c, i_{\sigma(d)}} \prod_{c \in C \cap B, d \sim_\pi c} J_{i_c, i_d} J_{i_{\sigma(c)}, i_d}. \end{aligned}$$

Each term of the sum is equal to 1 when $i_{\pi(k)} = i_k + N = i_{\eta(k)} \bmod(2N)$ and 0 otherwise. The above sum is therefore $(2N)^{\#(\pi \vee \eta)}$. Recalling that $\varepsilon(\sigma) = n - \#(\pi \vee \eta)$ yields the claim. The second point is easier to prove and we leave it to the reader. $\square$




Figure 1: In the following picture $\pi = \{\{1, 4\}, \{2, 3\}, \{5, 6\}, \{7, 10\}, \{8, 9\}, \{11, 12\}\}$, $\eta = \{\{1, 3\}, \{2, 5\}, \{4, 6\}, \{7, 8\}, \{9, 10\}, \{11, 12\}\}$, points of A are black, points of C are circled.

For any $z \in \mathbb{C}$ and any $\pi, \eta \in \mathcal{M}(2n)$, we set

$$G_{\pi, \eta}(z) = z^{\#(\pi \vee \eta)}.$$

For $|z| \geq 2n$, this matrix is invertible¹, let us denote by $W(z)$ its inverse. Equation (3) and Lemma 2.2 yield the following: for $I = (i_1, \dots, i_{2n}), J = (j_1, \dots, j_{2n}) \in \{1, \dots, 2N\}^{2n}$,

$$\int_{\text{USp}(2N)} S_{i_1, j_1} S_{i_2, j_2} \cdots S_{i_{2n}, j_{2n}} dS = \sum_{\pi, \eta \in \mathcal{M}(2n)} \varepsilon(\sigma_{\pi} \sigma_{\eta}) (-1)^{\text{inv}_2(\sigma_{\pi}, I) + \text{inv}_2(\sigma_{\eta}, J) + n} W_{\pi, \eta}(-2N), \quad (4)$$

where the sum is over all the partitions π, η such that $i_k = i_l + N \bmod(2N)$ (resp. $j_k = j_l + N \bmod(2N)$) as soon as $k \sim_{\pi} l$ (resp. $k \sim_{\eta} l$). Let us recall the two formulas when the group is the compact orthogonal group and the unitary group (see [2] for a proof). Let us define $\mathcal{M}(n, n) = \{\pi \in \mathcal{M}(2n) : \forall i, j \in \{1, \dots, n\}, i \not\sim_{\pi} j\}$ and let $\text{Wg}(z)$ be a pseudoinverse of $(G_{\pi, \eta}(z))_{\pi, \eta \in \mathcal{M}(n, n)}$. For any $I = (i_1, \dots, i_{2n}), J = (j_1, \dots, j_{2n}) \in \{1, \dots, N\}^{2n}$,

$$\int_{\text{O}(N)} O_{i_1, j_1} O_{i_2, j_2} \cdots O_{i_{2n}, j_{2n}} dO = \sum_{\pi, \eta \in \mathcal{M}(2n)} W_{\pi, \eta}(N), \quad (5)$$

where the sum is over all the partitions π, η such that $i_k = i_l$ (resp. $j_k = j_l$) as soon as $k \sim_{\pi} l$ (resp. $k \sim_{\eta} l$) and

$$\int_{\text{U}(N)} U_{i_1, j_1} U_{i_2, j_2} \cdots U_{i_n, j_n} \overline{U}_{i_{n+1}, j_{n+1}} \overline{U}_{i_{n+2}, j_{n+2}} \cdots \overline{U}_{i_{2n}, j_{2n}} dU = \sum_{\pi, \eta \in \mathcal{M}(n, n)} \text{Wg}_{\pi, \eta}(N), \quad (6)$$

where the sum is over all the partitions $\pi, \eta \in \mathcal{M}(n, n)$ such that $i_k = i_l$ (resp. $j_k = j_l$) as soon as $k \sim_{\pi} l$ (resp. $k \sim_{\eta} l$).

¹ We shall get an expression of the matrix $G(z)$ in terms of Jucy-Murphys elements, which shows that $G(z)$ is diagonalizable so that $G(z)$ always has a pseudo-inverse.

3 Reformulation in the Brauer algebra.

Let us remark that the endomorphism $\int_G g^n \mu(dg)$ of $V^{\otimes n}$ commutes with the action of G as soon as the measure μ is invariant by adjunction. We shall specify the element of $\text{End}_G(V^{\otimes n})$ when μ is Haar measure or the law of the Brownian motion. To begin with, let us describe the algebra $\text{End}_G(V^{\otimes n})$ as G belongs to the classical series.

3.1 Brauer algebra

Let us write I_n for the vector space with a basis indexed by $\mathcal{M}(2n)$. We give I_n the structure of an algebra as follows. For every $\pi \in \mathcal{M}(2n)$, let $\pi^+ = \{\{1\}, \dots, \{n\}\} \cup (\pi + n)$ and $\pi^- = \pi \cup \{\{2n+1\}, \dots, \{3n\}\} \in \mathcal{M}(3n)$. For any π, η , we denote by $b(\pi, \eta)$ the number of blocks of $(\pi_+ \vee \eta_-)$ included in $\{n+1, \dots, 2n\}$ and $\pi \circ \eta \in \mathcal{M}(2n)$ the partition obtained from $(\pi_+ \vee \eta_-)_{|\{1, \dots, n\} \cup \{2n+1, \dots, 3n\}}$ by shifting by $\{2n+1, \dots, 3n\}$ to $\{n+1, \dots, 2n\}$. We define

$$\pi \cdot \eta = z^{b(\pi, \eta)} \pi \circ \eta.$$

Lemma 3.1 ([6] Thm 4.4). *For any $z \in \mathbb{C}$, the bilinear map $(\pi, \eta) \in I_n^2 \mapsto \pi \cdot \eta$ endows I_n with a structure of an associative unitary algebra. We denote this algebra $\mathcal{B}_n(z)$.*

For any $\pi \in \mathcal{M}(2n)$, we shall represent the element $\pi \in \mathcal{B}_n(z)$ as a diagram with points of $\{1, \dots, 2n\}$ set on two lines as in the figure 2, so that multiplication in $\mathcal{B}_n(z)$ amounts to concatenation of diagram weighted by z powered to the number of inside loops. For any integers




Figure 2: Multiplication $\pi \cdot \eta$ with $\pi = \{\{1, 2\}, \{3, 4\}, \{5, 14\}, \{6, 7\}, \{8, 9\}, \{10, 11\}, \{12, 13\}\}$ and $\eta = \{\{1, 3\}, \{25\}, \{4, 6\}, \{7, 12\}, \{8, 11\}, \{9, 10\}, \{13, 14\}\}$.

$1 \leq a < b \leq n$, set

$$\tau_{a,b} = \{\{a, b\}, \{a+n, b+n\}\} \cup \{\{k, k+n\}, k \neq a, b\}$$

and

$$s_{a,b} = \{\{a, b+n\}, \{b, a+n\}\} \cup \{\{k, k+n\}, k \neq a, b\}.$$

This two kind of partitions generates $\mathcal{B}_n(z)$ as an algebra.


 Figure 3: The two elements $\tau_{a,b}$ et $s_{a,b}$ of $\mathcal{B}_n(z)$.

Lemma 3.2 ([5] § 7). *The algebra $\mathcal{B}_n(z)$ is isomorphic to the quotient of the free unital associative algebra on $\{\bar{\tau}_{a,b}, \bar{s}_{a,b} : 1 \leq a < b \leq n\}$ by the two-sided ideal generated by the following relations: for disctints integers a, b, c, d between 1 and n ,*

- (i) $\bar{\tau}_{a,b}^2 = z\bar{\tau}_{a,b}$, (ii) $\bar{s}_{a,b}^2 = 1$,
- (iii) $\bar{\tau}_{a,b}\bar{\tau}_{c,d} = \bar{\tau}_{c,d}\bar{\tau}_{a,b}$, (iv) $\bar{s}_{a,b}\bar{s}_{c,d} = \bar{s}_{c,d}\bar{s}_{a,b}$,
- (v) $\bar{s}_{c,d}\bar{\tau}_{a,b}\bar{s}_{c,d} = \bar{\tau}_{(c\ d).a,(c\ d).b}$, (vi) $\bar{s}_{a,b}\bar{s}_{b,c}\bar{s}_{a,b} = \bar{s}_{b,c}\bar{s}_{a,b}\bar{s}_{b,c}$,
- (vii) $\bar{\tau}_{a,b}\bar{\tau}_{b,c} = \bar{s}_{a,c}\bar{\tau}_{b,c}$.

Let us describe the standard representations of $\mathcal{B}_n(N)$ and $\mathcal{B}_n(-2N)$. For any $\pi \in \mathcal{M}(2n)$, let us define the element of $\text{End}(V^{\otimes n})$

$$\rho_O(\pi) = \sum E_{i_{n+1}, i_1} \otimes \cdots \otimes E_{i_{2n}, i_n},$$

the sum being over all $i_1, \dots, i_{2n} \in \{1, \dots, N\}$ such that $i_k = i_l$ for any k, l such that $k \sim_\pi l$. For any $1 \leq a < b \leq n$, let us denote by $\langle a\ b \rangle = \rho_O(\tau_{a,b})$ and $(a\ b) = \rho_O(s_{a,b})$.

Lemma 3.3. *The application ρ_O extends linearly to an algebra homomorphism from $\mathcal{B}_n(N)$ to $\text{End}(V^{\otimes n})$.*

For any $1 \leq a < b \leq n$, let

$$\langle a\ b \rangle_\omega = \sum_{1 \leq i_a, i_{a+n}, i_b, i_{b+n} \leq 2N} J_{i_a, i_b} J_{i_{a+n}, i_{b+n}} \text{Id}^{\otimes a-1} \otimes E_{i_{a+n}, i_a} \otimes \text{Id}^{\otimes b-a-1} \otimes E_{i_{b+n}, i_b} \otimes \text{Id}^{\otimes n-b},$$

$$\rho_S(s_{a,b}) = -(a\ b) \in \text{End}(V^{\otimes n}) \text{ and } \rho_S(\tau) = -\langle a\ b \rangle_\omega.$$

Lemma 3.4. *The application ρ_S extends to an algebra homomorphism from $\mathcal{B}_n(-2N)$ to $\text{End}(V_{\mathbb{C}}^{\otimes n})$.*

Proof. One checks that the seven relations given in Lemma 3.2 are satisfied. $\square$

3.2 Expectation of tensors as elements of the Brauer algebra.

Let us fix a family of permutations $(\sigma_\pi)_{\pi \in \mathcal{M}(2n)}$ such that $\sigma_\pi \pi_0 = \pi$. For every $\sigma \in \mathfrak{S}_{2n}$, define R_σ as the endomorphism of I_n such that for any $\pi \in \mathcal{M}(2n)$,

$$R_\sigma(\pi) = \sigma_\pi \sigma \pi_0.$$

For any $x \in \mathbb{C}[\mathfrak{S}_{2n}]$, let us define R_x by extending the previous definition linearly. Let us point out that this does not define a right action of $\mathfrak{S}_{2n}$. Set

$$G(z) = \sum_{\pi \in \mathcal{M}(2n)} z^{\#(\pi \vee \pi_0)} \sigma_\pi \in \mathbb{C}[\mathfrak{S}_{2n}].$$

Lemma 3.5. *The matrix of the endomorphism $R_{G(z)}$ of I_n in its canonical basis is $(G_{\mu,\nu})_{\mu,\nu \in \mathcal{M}(2n)}$.*

Proof. For any $\pi, \eta, \mu \in \mathcal{M}(2n)$ with $R_{\sigma_\pi}(\eta) = \mu$, $\nu \vee \mu = \sigma_\eta(\pi_0 \vee \pi)$ and $\#(\pi \vee \pi_0) = \#(\mu \vee \nu)$. $\square$

Let us point out the following fact proved in [10].

Proposition 3.6 ([10], Proposition 3). *There exists a family of permutations $(\sigma_\pi)_{\pi \in \mathcal{M}(2n)}$ such that $\sigma_\pi \pi_0 = \pi$ for any $\pi \in \mathcal{M}(2n)$ and*

$$G(z) = \prod_{k=1}^n (z + X_{2k-1}).$$

The knowledge of the spectrum of Jucy-Murphy elements yields that whenever $z \notin \{-2n+2, -2n+3, \dots, 2n-3, 2n-2\}$, $G(z) \in \mathbb{C}[\mathfrak{S}_{2n}]$ is invertible.

Lemma 3.7. *i) For any $h \in \mathcal{H}_{2n}$, $h.G(z)\pi_0 = G(z)\pi_0$, and $h.G(z)^{-1}\pi_0 = G(z)^{-1}\pi_0$.*

ii) For any $z \notin \{-2n+2, -2n+3, \dots, 2n-3, 2n-2\}$, $R_{G(z)}$ is invertible with inverse $R_{G(z)^{-1}}$.

Proof. i) Let us write $\mathcal{P}_{\mathcal{H}_{2n}} = \frac{1}{|\mathcal{H}_{2n}|} \sum_{h \in \mathcal{H}_{2n}} h \in \mathbb{C}[\mathfrak{S}_{2n}]$. The two equalities are equivalent to $G(z)\mathcal{P}_{\mathcal{H}} = \mathcal{P}_{\mathcal{H}}G(z)$. Moreover, for any $h \in \mathcal{H}_{2n}$ and $\pi \in \mathcal{M}(2n)$, if $h\sigma_\pi = \sigma_{\pi'}h'$ with $\pi' \in \mathcal{M}(2n)$, $h' \in \mathcal{H}_{2n}$, then $\#\pi \vee \pi_0 = \#\pi' \vee \pi_0$. Therefore for any $h \in \mathcal{H}_{2n}$, $hG(z)\mathcal{P}_{\mathcal{H}_{2n}} = G(z)\mathcal{P}_{\mathcal{H}_{2n}}$.

ii) Let us notice that I_n is isomorphic to $\mathbb{C}[\mathfrak{S}_{2n}].\mathcal{P}_{\mathcal{H}_{2n}}$ and thanks to point i), multiplication on the right by $P(G(z))$ for any polynomial P is conjugate by this isomorphism to $R_{P(G(z))}$. Hence $R_{G(z)}$ has as a pseudoinverse $R_{G(z)^{-1}}$. $\square$

We can now give another formulation of the results of the last section. For any $r, s, n \in \mathbb{N}$ such that $r \leq s \leq n$ and $s-r$ is even, let us denote $\tau_{[r,s]} = \prod_{1 \leq 2i \leq s-r} \tau_{r+2i-1, r+2i} \in \mathcal{B}_n(z)$.

Proposition 3.8. *Let $\mathcal{I}_{2n}(z) = \frac{1}{2^n n!} \sum_{\sigma \in \mathfrak{S}_{2n}} \sigma G(z)^{-1} \tau_{[1,2n]} \sigma^{-1}$,*

$$\int_{\text{USp}(2N)} S^{\otimes 2n} dS = \rho_S(\mathcal{I}_{2n}(-2N))$$

and

$$\int_{\text{O}(N)} O^{\otimes 2n} dO = \rho_O(\mathcal{I}_{2n}(N)).$$

Proof. Let us prove the first formula for the symplectic group. Let us observe that for every $\pi \in \mathcal{M}(2n)$, the following equality holds in $\text{End}(V^{\otimes 2n})$,

$$\langle w_\eta, \cdot \rangle w_\pi = (-1)^n \rho_S(\sigma_\pi \tau_{[1,2n]} \sigma_\eta).$$

What is more, Lemma 2.2 implies

$$\langle w_\pi, w_\eta \rangle = (-1)^n G_{\pi,\eta}(-2N).$$

Let us extend by linearity the definition of w . to I_n . The second assertion of Lemma 3.7 implies

$$\begin{aligned} \sum_{\pi, \eta \in \mathcal{M}(2n)} (-1)^n W_{\pi,\eta}(-2N) \langle w_\eta, \cdot \rangle w_\pi &= (-1)^n \sum_{\eta \in \mathcal{M}(2n)} \langle w_\eta, \cdot \rangle w_{R_{G(-2N)^{-1}}(\eta)} \\ &= \sum_{\eta \in \mathcal{M}(2n)} \rho_S(\sigma_\eta G(-2N)^{-1} \tau_{[1,2n]} \sigma_\eta^{-1}). \end{aligned}$$

The first point of Lemma 3.7 yields the result. $\square$

We shall now give a statement of our main result which is a generalization of Proposition 3.8 when the measure is not the Haar measure but the law of a Brownian motion at a fixed time. For any $x \in \mathfrak{g}$, let $\mathcal{L}_x$ be the left invariant first order differential operator defined by setting for any differentiable function f and any $g \in G$,

$$\mathcal{L}_x f(g) = \frac{d}{dt} f(g \exp(tx))|_{t=0}.$$

As G is any of the three groups $O(N)$, $USp(N)$ and $U(N)$, we write V for the fundamental representation and $\langle \cdot, \cdot \rangle$ for the scalar product on $\mathfrak{g}$ given for any $x, y \in \mathfrak{g}$ by

$$\langle x, y \rangle = -\dim(V) \text{Tr}_V(xy).$$

Let $(x_i)_{1 \leq i \leq d}$ be an orthonormal basis of $\mathfrak{g}$ and define $\Delta_G = \sum_{i=1}^d \mathcal{L}_{x_i} \circ \mathcal{L}_{x_i}$. The operator Δ_G is independent of the orthonormal basis $(x_i)_{1 \leq i \leq d}$ and is called the Laplacian on G . The Brownian motion on G is the diffusion with generator $\frac{1}{2} \Delta_G$. We shall write $(O_t)_{t \geq 0}$ (resp. $(S_t)_{t \geq 0}$ and $(U_t)_{t \geq 0}$) for the Brownian motion on $SO(N)$ (resp. on $USp(N)$ and on $U(N)$) issued from the identity.

For any $C \subset \{1, \dots, n\}$, set

$$Z_C(z) = \frac{(1 - z^{-1})|C|}{2} + z^{-1} \sum_{a < b: a, b \in C} s_{a,b} \in \mathcal{B}_n(z)$$

and for $1 \leq i \leq n$,

$$Z_i = Z_{\{1, \dots, i\}}(z).$$

For $t \in \mathbb{R}$ and $k \in \mathbb{N}$, let $s_t^{(k)}$ and $g_t^{(k)}$ be the holomorphic symmetric functions defined by:

$$s_t^{(k)}(z_1, \dots, z_k) = \prod_{1 \leq i < j \leq k} (z_j - z_i)^{-1} \det \begin{pmatrix} z_1^{k-1} & z_1^{k-2} & \dots & z_1 & e^{-tz_1} \\ z_2^{k-1} & z_2^{k-2} & \dots & z_2 & e^{-tz_2} \\ \vdots & \vdots & & \vdots & \vdots \\ z_k^{k-1} & z_k^{k-2} & \dots & z_k & e^{-tz_k} \end{pmatrix}$$

and

$$g_t^{(k)}(z_1, \dots, z_k) = -z_1^{-1} \dots z_k^{-1} s_t^{(k)}(z_1, \dots, z_k).$$

For any integers n and p such that $0 < 2p < n$, let

$$\mathcal{I}_{n,t}^p(z) = \frac{1}{2^p z^p (n-2p)!} \sum_{\sigma \in \mathfrak{S}_n} \sigma g_t^{(p+1)}(Z_{n-2p}, Z_{n-2p+2}, \dots, Z_n) \tau_{[n-2p+1, n]} \sigma^{-1}.$$

If $n = 2p$, set

$$\mathcal{I}_{n,t}^p(z) = \frac{1}{2^p p!} \sum_{\sigma \in \mathfrak{S}_n} \sigma G(z)^{-1} \left(1 - s_t^{(p)}(1, 2(1 + z^{-1}X_3), \dots, p(1 + z^{-1}X_{2p-1})) \right) \tau_{[1, n]} \sigma^{-1}.$$

Let us write $\mathcal{I}_{n,t}^0(z) = e^{-tZ_n}$ and $\mathcal{I}_{n,t}(z) = \sum_{0 \leq 2p \leq n} \mathcal{I}_{n,t}^p(z)$. Our main result is the following formulas.

Theorem 3.9. *For any $t \in \mathbb{R}_+$ and $n \in \mathbb{N}$,*

$$\mathbb{E}[O_t^{\otimes n}] = \rho_O(\mathcal{I}_{n,t}(N))$$

and

$$\mathbb{E}[S_t^{\otimes n}] = \rho_S(\mathcal{I}_{n,t}(-2N)).$$

3.3 Walled Brauer algebra

For two integers n and m , let $\mathcal{M}(n, m)$ be the set of partitions which does not connect $\{1, \dots, n\}$ with $\{2n + m + 1, \dots, 2(n + m)\}$ nor $\{n + 1, \dots, n + m\}$ with $\{n + m + 1, \dots, 2n + m\}$. We note $\mathcal{B}_{n,m}(z)$ the vector space spanned by $\mathcal{M}(n, m)$. Let us notice that $\mathcal{B}_{n,m}(z)$ is a subalgebra of $\mathcal{B}_{n+m}(z)$.




Figure 4: An element of $\mathcal{B}_{n,m}(z)$.

Let us denote by $\tilde{\pi}_0 \in \mathcal{M}(n, n)$ the partition $\{1, n + 1\}, \dots, \{n, 2n\}$. For any $\pi \in \mathcal{M}(n, n)$, let $\tilde{\sigma}_\pi \in \mathfrak{S}_n$ be the permutation such that $\tilde{\sigma}_\pi(\tilde{\pi}_0) = \pi$. For any permutation $\sigma \in \mathfrak{S}_n$, let us write $\#\sigma$ for the number of cycles of σ and set

$$\Omega(z) = \sum_{\sigma \in \mathfrak{S}_n} z^{\#\sigma} \sigma \in \mathbb{C}[\mathfrak{S}_n].$$

For any $\pi, \eta \in \mathcal{M}(n, n)$, let us observe that $\#\sigma_\pi^{-1}\sigma_\eta = \#(\sigma_\pi^{-1}\sigma_\eta \times \text{Id}_n(\tilde{\pi}_0) \vee \tilde{\pi}_0) = \#\pi \vee \eta$, hence

$$G(z)_{\pi, \eta} = \Omega(z)(\sigma_\pi^{-1}\sigma_\eta)$$

and

$$\text{Wg}(z)_{\pi, \eta} = \text{Wg}(z)(\sigma_\pi^{-1}\sigma_\eta).$$

The following formula is proved in [10].

Proposition 3.10 ([10], Prop. 1). *For any $z \in \mathbb{C}$,*

$$\Omega(z) = \prod_{i=1}^n (z + X_i).$$

For $r, s, n, m \in \mathbb{N}$, such that $1 \leq r \leq s \leq n \leq m$, let

$$\tilde{\tau}_{[r,s]} = \prod_{r \leq k \leq s} \tau_{k, k+n}.$$

We can now give another formulation of expression (6).

Proposition 3.11. *Let $\mathcal{I}_{n,n}(z) = \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n \times \mathfrak{S}_n} \sigma \Omega(z)^{-1} \tilde{\tau}_{[1,n]} \sigma^{-1} \in \mathcal{B}_{n,n}(z)$,*

$$\int_{U(N)} U^{\otimes n} \otimes \bar{U}^{\otimes n} dU = \rho_O(\mathcal{I}_{n,n}(N)).$$

We shall state the counterpart of Proposition 3.11 for the Brownian motion. Let n, m be two integers such that $n \leq m$. Let us denote by p^w the projection from $\mathcal{B}_{n+m}(z)$ to $\mathcal{B}_{n,m}(z)$ such that $p^w(\pi) = 0$ for any $\pi \in \mathcal{M}(2(n+m)) \setminus \mathcal{M}(n, m)$. For any $C \subset \{1, \dots, n+m\}$, set $Y_C = p^w(Z_C) + \frac{|C|z^{-1}}{2} \in \mathcal{B}_{n,m}(z)$ and for any $i \leq n$, set

$$Y_i = Y_{\{1, \dots, i\} \cup \{n+1, \dots, m+i\}}.$$

For any integer p such that $1 \leq p < n$, set

$$\mathcal{I}_{n,m,t}^p(z) = \frac{1}{z^p(n-p)!(m-p)!} \sum_{\sigma \in \mathfrak{S}_n \times \mathfrak{S}_m} \sigma g_t^{(p+1)}(Y_{n-p}, Y_{n-p+1}, \dots, Y_n) \tilde{\tau}_{[n-p+1,n]} \sigma^{-1}.$$

For $n < m$, set

$$\mathcal{I}_{n,m,t}^n(z) = \frac{1}{z^n(m-p)!} \sum_{\sigma \in \mathfrak{S}_n \times \mathfrak{S}_m} \sigma g_t^{(p+1)}(Y_0, Y_1, \dots, Y_n) \tilde{\tau}_{[1,n]} \sigma^{-1}$$

and for $n = m$,

$$\mathcal{I}_{n,n,t}^n(z) = \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n \times \mathfrak{S}_n} \sigma \Omega(z)^{-1} \left(1 - s_t^{(p)}(1, 2(1 + z^{-1}X_2), \dots, n(1 + z^{-1}X_n)) \right) \tilde{\tau}_{[1,n]} \sigma^{-1}.$$

Let us write $\mathcal{I}_{n,m,t}^0(z) = e^{-tY_n}$ and $\mathcal{I}_{n,m,t}(z) = \sum_{p=0}^n \mathcal{I}_{n,m,t}^p(z)$. The formula of Proposition 3.11 has the following deformation.

Theorem 3.9. *For any nonnegative integers $n \leq m$ and $t \geq 0$,*

$$\mathbb{E}[U_t^{\otimes n} \otimes \bar{U}_t^{\otimes m}] = \rho_O(\mathcal{I}_{n,m,t}(N)).$$

3.4 Limit as time goes to infinity

For any compact connected Lie group G , the law at a time t of the Brownian motion issued from the identity converges to the Haar measure as t goes to infinity. Hence formulas of Theorem 3.9 yields formulas for the Haar measure. Let us check that our formulas yield the one of B. Collins and P. Śniady. Observe that $O(N)$ is not connected and that the law of a Brownian motion at time t issued from the identity converges to the Haar measure on $SO(N)$ as t goes to infinity. The following lemma shows that the integration of all homogeneous polynomials but the determinant with respect to the latter equals integration with respect to the Haar measure on $O(N)$. Let us denote by ε_n the idempotent $\frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \varepsilon(\sigma) \sigma \in \mathbb{C}[\mathfrak{S}_n]$.

Lemma 3.12. *Let n, m and N be fixed integers with $n \leq m$. Let us set $\mathcal{I}_n(z) = 0$ if n is odd and $\mathcal{I}_{n,m}(z) = 0$ if $n < m$. As t goes to infinity,*

$$\begin{aligned} \rho_O(\mathcal{I}_{n,t}(N)) &\longrightarrow \rho_O(\mathcal{I}_n(N) + 1_{\{n=N\}} \varepsilon_N), \\ \rho_S(\mathcal{I}_{n,t}(-2N)) &\longrightarrow \rho_S(\mathcal{I}_n(-2N)) \end{aligned}$$

and

$$\rho_O(\mathcal{I}_{n,m,t}(N)) \longrightarrow \rho_O(\mathcal{I}_{n,m}(N)).$$

Proof. Using Theorem 3.9, Lemma 4.5 and assertion iii) of Lemma 4.3, it is sufficient to prove that the spectrum of the homomorphisms $\rho_O(Z_i(N))$, $\rho_S(Z_i(-2N))$ and $\rho_O(Y_i(N))$ are strictly positive for all $i \leq n$. The operator norm of a transposition is less than one, hence for any $x > 0$ and $i, j \in \mathbb{N}$ with $i > j$, the spectrum of $Z_i(x)$ and $\rho_S(Z_i(-x))$, resp. $Y_i(x)$ is bounded below by $\frac{1}{2|x|}((|x| - 1)i - i(i - 1)) = \frac{i}{2|x|}(|x| - i)$, resp. $\frac{1}{2|x|}(i(|x| - i + 1) + (m - n + i)(|x| - m + n - i + 1))$. Therefore the conclusion of the Lemma follows for N large enough. To prove the Lemma for small N , notice that

$$zZ_i(z) = \sum_{k=1}^i \left(\frac{z-1}{2} + X_k \right)$$

and

$$zY_i(z) = \sum_{k=1}^i \left(\frac{z}{2} + X_k \right) + \alpha \sum_{k=1}^{m-n+i} \left(\frac{z}{2} + X_k \right) \alpha^{-1},$$

where α is an involution of $\{1, \dots, m+n\}$ such that for any $k \leq m$, $\alpha(k) = k+n$. Observe that the two terms of the RHS of the last equality commute and can be codiagonalized. The explicit knowledge of the spectrum of Jucy-Murphy elements (see [9]) and the decomposition of $V^{\otimes n}$ into irreducible $\mathfrak{S}_n$ components (see [2] or [7]) allow to conclude in all the three cases but the first one as $n = N$. In that case, denote by L the line of $V^{\otimes N}$ spanned by the determinant and $L^\perp$ its orthogonal in $V^{\otimes N}$. The previous analysis shows that $\mathbb{E}[O_t^{\otimes N}]$ restricted to $L^\perp$ goes to zero as t goes to infinity. What is more, $\mathbb{E}[O_t^{\otimes N}]$ acts trivially on L . $\square$

4 Expectation of tensors under the Brownian motion

The rest of the paper is devoted to the proof of our main result Theorem 3.9.

4.1 The Casimir element and its representations

Let G be a compact Lie group with Lie algebra $\mathfrak{g}$ of dimension d with a G -invariant scalar product $\langle \cdot, \cdot \rangle$. For any orthonormal basis $(x_i)_{1 \leq i \leq d}$ of $\mathfrak{g}$ with respect to the scalar product $\langle \cdot, \cdot \rangle$, let $c_{\mathfrak{g}} = \sum_{i=1}^d x_i^2 \in \mathcal{U}(\mathfrak{g})$. The element $c_{\mathfrak{g}}$ does not depend on the orthonormal basis $(x_i)_{1 \leq i \leq d}$ and is called the Casimir element. The application $\mathcal{L}$ extends to an isomorphism of associative algebra between $\mathcal{U}(\mathfrak{g})$ and the algebra of left-invariant differential operators on G which sends $c_{\mathfrak{g}}$ to Δ_G . Let $(g_t)_{t \geq 0}$ be a Brownian motion issued from Id. For any finite dimensional representation (V_ρ, ρ) and any $t \geq 0$,

$$\frac{d}{dt} \mathbb{E}[\rho(g_t)] = \mathbb{E}\left[\frac{1}{2} \Delta_G(\rho)(g_t)\right] = \mathbb{E}\left[\frac{1}{2} \rho(c_{\mathfrak{g}}) \rho(g_t)\right] \in \text{End}(V_\rho)$$

and

$$\mathbb{E}[\rho(g_t)] = \exp\left(\frac{t}{2} \rho(c_{\mathfrak{g}})\right).$$

Let us observe that the element $c_{\mathfrak{g}}$ is invariant by adjunction, hence $\rho(c_{\mathfrak{g}}) \in \text{End}_G(V_\rho)$. Our aim here is to specify $\exp(\frac{t}{2} \rho(c_{\mathfrak{g}}))$ as an element of $\text{End}_G(V_\rho)$ when (V_ρ, ρ) is a tensor power of the fundamental representation of a classical group. All the following formulas but the last one are proved in Proposition 2.6 and 2.8 of [7].

Lemma 4.1. *For any integers n and m , let us write*

$$\Delta_{\mathcal{B}_n}(z) = -\frac{(1-z^{-1})n}{2} + z^{-1} \sum_{1 \leq a < b \leq n} \tau_{a,b} - s_{a,b} \in \mathcal{B}_n(z)$$

and

$$\Delta_{\mathcal{B}_{n,m}}(z) = p^w(\Delta_{\mathcal{B}_{n+m}}(z)) - \frac{n+m}{2z} \in \mathcal{B}_{n+m}(z).$$

The Casimir elements of the classical groups have the following tensor representations:

$$\rho_{V^{\otimes n}}(c_{\mathfrak{o}(N)}) = 2\rho_O(\Delta_{\mathcal{B}_n}(N)),$$

$$\rho_{V^{\otimes n}}(c_{\mathfrak{usp}(2N)}) = 2\rho_S(\Delta_{\mathcal{B}_n}(-2N))$$

and

$$\rho_{V^{\otimes n}} \otimes \bar{\rho}_{V^{\otimes m}}(c_{\mathfrak{u}(N)}) = 2\rho_O(\Delta_{\mathcal{B}_{n+m}}(N)).$$

Proof. Let us prove the third equality. Let us choose the orthonormal basis $\{\frac{1}{\sqrt{2N}}(E_{k,l} - E_{l,k}), \frac{i}{\sqrt{2N}}(E_{k,l} + E_{l,k}) : 1 \leq k < l \leq N\}$ of $(\mathfrak{u}(N), \langle \cdot, \cdot \rangle)$ and compute the Casimir as an element of the real algebra $\mathcal{U}(\mathfrak{gl}_N(\mathbb{C}))$:

$$Nc_{\mathfrak{u}(N)} = \frac{1}{2} \sum_{1 \leq k, l \leq N} (1 + i \otimes i) E_{k,l} \otimes E_{k,l} + (i \otimes i - 1) E_{k,l} \otimes E_{l,k} \in \mathcal{U}(\mathfrak{gl}_N(\mathbb{C})).$$

Considering $\rho_{V^{\otimes n}} \otimes \bar{\rho}_{V^{\otimes m}}$ as a representation of the real algebra $\mathcal{U}(\mathfrak{gl}_N(\mathbb{C}))$ implies that

$$\begin{aligned} N\rho_{V^{\otimes n}} \otimes \bar{\rho}_{V^{\otimes m}}(c_{\mathfrak{u}(N)}) &= (n+m)N\text{Id}_{V^{\otimes n+m}} + 2 \sum_{1 \leq a \leq n < b \leq n+m} \langle a \ b \rangle - 2 \sum_{\substack{1 \leq a < b \leq n+m \\ a, b \leq n \text{ or } a, b > n}} (a \ b) \\ &= \rho_O \left((n+m)N\text{Id}_{\mathcal{B}_{n+m}(N)} + 2p^w \left(\sum_{1 \leq a < b \leq n+m} \tau_{a,b} - s_{a,b} \right) \right). \end{aligned}$$

Dividing by N yields the result.

Let us check the formula for the symplectic group. It is proved in Proposition 2.8 of [7] that

$$-2Nc_{\mathfrak{usp}(2N)} = - \sum_{1 \leq a, b, c, d \leq 2N} J_{a,c} J_{b,d} E_{a,b} \otimes E_{c,d} + \sum_{1 \leq a, b \leq 2N} E_{a,b} \otimes E_{b,a} \in \mathcal{U}(\mathfrak{gl}_N(\mathbb{C})).$$

Note that the scalar product $\langle \cdot, \cdot \rangle$ on $\mathfrak{usp}(2N)$ used in [7] is not normalized, the Casimir here is therefore $2N$ times larger. Considering $\rho_{V^{\otimes n}}$ as a representation of the algebra $\mathcal{U}(\mathfrak{gl}_N(\mathbb{C}))$ yields

$$\begin{aligned} -2N\rho_{V^{\otimes n}}(\Delta_{\mathfrak{usp}(2N)}) &= 2(2N+1)n + 2 \sum_{1 \leq a < b \leq n} (a \ b) - \langle a \ b \rangle_\omega \\ &= 2\rho_S \left((2N+1)n + \sum_{1 \leq a < b \leq n} \tau_{a,b} - s_{a,b} \right). \end{aligned}$$

Dividing by $-2N$ gives the announced formula. $\square$

4.2 A decomposition of the Brauer algebra

For any subset A of $\{1, \dots, n\}$, let us denote by $\mathcal{M}_A$ the set of $\pi \in \mathcal{M}(2n)$ such that $n + A$ is the maximal subset of $\{n + 1, \dots, 2n\}$ not connected to $\{1, \dots, 2n\} \setminus (A + n)$. For any integer p less than n , let $\mathcal{M}_p$ be the vector space spanned by $\cup_{A \subset \{1, \dots, n\}: |A|=2p} \mathcal{M}_A$ and $(P_{\mathcal{M}_p})_{0 \leq 2p \leq n}$ the family of projectors given by the decomposition $\mathcal{B}_n(z) = \bigoplus_{0 \leq 2p \leq n} \mathcal{M}_p$.

Lemma 4.2. *Let $n \leq m$ be two integers. Let $A \subset \{1, \dots, n\}$ and $\pi \in H_A$. For any $a \neq b \notin A$, $s_{a,b}\pi \in H_A$ and $\tau_{a,b}\pi \in H_{A \cup \{a,b\}}$. What is more,*

$$\Delta_{\mathcal{B}_n}(z)\pi = \left(-\frac{(1-z^{-1})(n-|A|)}{2} + z^{-1} \sum_{a < b: a, b \notin A} \tau_{a,b} - s_{a,b} \right) \pi$$

and if $\pi \in \mathcal{M}(n, m)$,

$$\Delta_{\mathcal{B}_{n,m}}(z)\pi = \left(-\frac{(n+m-|A|)}{2} + z^{-1} \sum_{a < b: a, b \notin A} p^w (\tau_{a,b} - s_{a,b}) \right) \pi.$$

Proof. We shall prove the two formulas and leave the first point to the reader. For any $k \in \{1, n\}$, we note $k' = k + n$ and $\pi(k') \in \{1, \dots, 2n\}$ for the element such that $\{k', \pi(k')\}$ is a block of π . Let $1 \leq k \neq l \leq n$, we compute $\tau_{k,l}\pi$ by taking into account whether k' or l' are in A and in the same block of π (see figure 5):

- a) If $k', l' \notin A$, $\tau_{k,l}\pi \in H_{A \cup \{k,l\}}$,
- b) if $k', l' \in A$ and $l' \neq \pi(k')$, $\tau_{k,l}\pi = s_{k, \pi(l')-n}\pi$,
- c) if $k', l' \in A$ and $l' = \pi(k')$, $\tau_{k,l}\pi = z s_{k,l}\pi = z\pi$,
- d) if $k' \in A$ and $l' \notin A$, $\tau_{k,l}\pi = s_{\pi(k')-n, l}\pi$.

We deduce that

$$\left(\sum_{1 \leq k < l \leq n} \tau_{k,l} - s_{k,l} \right) \pi = \left(\frac{(z-1)|A|}{2} + \sum_{1 \leq a < b \leq n: a, b \notin A} \tau_{a,b} - \sum_{1 \leq a < b \leq n: a, b \notin A} s_{a,b} \right) \pi \in \mathcal{B}_n(z).$$


Note that if $\pi \in \mathcal{M}(n, m)$, transpositions appearing in c) does not occur if we consider just elements in $\mathcal{B}_{n,m}(z)$ and

$$p^w \left(\sum_{1 \leq k < l \leq n} \tau_{k,l} - s_{k,l} \right) \pi = \left(\frac{z|A|}{2} + \sum_{1 \leq a < b \leq n: a, b \notin A-n} \tau_{a,b} - \sum_{1 \leq a < b \leq n: a, b \notin A-n} s_{a,b} \right) \pi \in \mathcal{B}_{n,m}(z).$$

The result follows from this two formulas. □

For any integers r and k , let us denote by

$$h_r(z_1, \dots, z_k) = \sum_{\substack{l_1, \dots, l_k \geq 0 \\ l_1 + \dots + l_k = r}} z_1^{l_1} \cdots z_k^{l_k},$$


 Figure 5: Left multiplication of an element of $\mathcal{M}_A$ by $\tau_{k,l}$.

the complete symmetric polynomial of degree r . We recall the following classical fact that complete polynomials agree with Schur functions of the partitions $(r, 0)$. A proof of the first point of the following lemma can be found in [8]. The two other assertions follow from it.

Lemma 4.3. *i) For any integers r and l ,*

$$h_r^{(l)}(z_1, \dots, z_l) = \prod_{1 \leq i < j \leq l} (z_j - z_i)^{-1} \det \begin{pmatrix} z_1^{r+l-1} & z_1^{l-2} & z_1^{l-3} & \cdots & 1 \\ z_2^{r+l-1} & z_2^{l-2} & z_2^{l-3} & \cdots & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ z_l^{r+l-1} & z_l^{l-2} & z_l^{l-3} & \cdots & 1 \end{pmatrix}.$$

ii) For any $t \in \mathbb{R}$,

$$\sum_{r \geq 0} \frac{(-t)^{r+l}}{(r+l)!} h_r^{(l)}(z_1, \dots, z_l) = (-1)^l z_1^{-1} \cdots z_l^{-1} \left(1 - s_t^{(l)}(z_1, \dots, z_l) \right)$$

and

$$\sum_{r \geq 0} \frac{(-t)^{r+l}}{(r+l)!} h_r^{(l+1)}(z_1, \dots, z_{l+1}) = (-1)^{l+1} z_1^{-1} \cdots z_{l+1}^{-1} s_t^{(l+1)}(z_1, \dots, z_{l+1}).$$

iii) For any distinct complex numbers $z_1, \dots, z_l$,

$$z_1^{-s} z_2^{-1} \cdots z_l^{-1} s_t^{(s+l-1)}(z_1, \dots, z_1, z_2, \dots, z_l) = \frac{1}{(s-1)!} \frac{\partial^{s-1}}{\partial z_1^{s-1}} z_1^{-1} \cdots z_l^{-1} s_t^{(l)}(z_1, z_2, \dots, z_l).$$

The following lemma is the main computation in the proof of Theorem 3.9.

Lemma 4.4. *i) For any $k < p$, $P_{\mathcal{M}_p} \Delta_{\mathcal{B}_n}^k = P_{\mathcal{M}_p} \Delta_{\mathcal{B}_{n,m}}^k = 0$.*

ii) For $k \geq p$, if $2p < n$,

$$z^p P_{\mathcal{M}_p} \Delta_{\mathcal{B}_n}^k = \frac{(-1)^{k-p}}{2^p(n-2p)!} \sum_{\sigma \in \mathfrak{S}_n} \sigma h_{k-p}^{(p+1)}(Z_{n-2p}, \dots, Z_n) \tau_{[n-2p+1, n]} \sigma^{-1}$$

and if $p < n$ or $n < m$,

$$z^p P_{\mathcal{M}_p} \Delta_{\mathcal{B}_{n,m}}^k = \frac{(-1)^{k-p}}{(n-p)!(m-p)!} \sum_{\sigma \in \mathfrak{S}_n \times \mathfrak{S}_m} \sigma h_{k-p}^{(p+1)}(Y_{n-p}, \dots, Y_n) \tilde{\tau}_{[n-p+1, n]} \sigma^{-1}.$$

iii) For $k \geq p$, if $n = 2p$,

$$z^p P_{\mathcal{M}_p} \Delta_{\mathcal{B}_n}^k = \frac{(-1)^{k-p}}{2^p} \sum_{\sigma \in \mathfrak{S}_n} \sigma h_{k-p}^{(p)}(Z_1, \dots, Z_n) \tau_{[1, n]} \sigma^{-1}$$

and if $n = p$ and $n = m$

$$z^p P_{\mathcal{M}_p} \Delta_{\mathcal{B}_{n,n}}^k = (-1)^{k-p} \sum_{\sigma \in \mathfrak{S}_n \times \mathfrak{S}_n} \sigma h_{k-n}^{(n)}(Y_1, \dots, Y_n) \tilde{\tau}_{[1, n]} \sigma^{-1}.$$

Proof. Using the assertion of Lemma 4.2, we deduce that for $0 \leq k < p$, $\Delta_{\mathcal{B}_n(z)}^k \in \bigoplus_{0 \leq 2p \leq 2k} \mathcal{M}_p$, which implies the point i). We shall prove points ii) and iii) only for $\Delta_{\mathcal{B}_n(z)}$, the same proof applies to the computation of powers of $\Delta_{\mathcal{B}_{n,m}}(z)$. For $q_1 < q_2, \dots, q_{2p-1} < q_{2p}$ distinct non-zero integers smaller than n , note $B = \{1, \dots, n\} \setminus \{q_1, q_2, \dots, q_{2p}\}$ and for $k \geq p$, let $F_k(q_1, q_2, \dots, q_{2p-1}, q_{2p}) \in \mathcal{B}_n(z)$ be the element

$$\sum_{\substack{l_0, l_1, \dots, l_p \geq 0 \\ l_0 + \dots + l_p = k-p}} Z_B^{l_0} \tau_{q_1, q_2} Z_{B \cup \{q_1, q_2\}}^{l_1} \tau_{q_3, q_4} \dots Z_{\{1, \dots, n\} \setminus \{q_{2p-1}, q_{2p}\}}^{l_{p-1}} \tau_{q_{2p-1}, q_{2p}} Z_{\{1, \dots, n\}}^{l_p},$$

if $B \neq \emptyset$, and

$$\sum_{\substack{l_1, \dots, l_p \geq 0 \\ l_1 + \dots + l_p = k-p}} \tau_{q_1, q_2} Z_{\{q_1, q_2\}}^{l_1} \tau_{q_3, q_4} \dots Z_{\{1, \dots, n\} \setminus \{q_{2p-1}, q_{2p}\}}^{l_{p-1}} \tau_{q_{2p-1}, q_{2p}} Z_{\{1, \dots, n\}}^{l_p},$$

otherwise. For $k \geq p$, Lemma 4.2 yields

$$z^p P_{\mathcal{M}_p} \Delta_{\mathcal{B}_n}^k(z) = (-1)^{k-p} \sum_{\substack{q_1 < q_2, \dots, q_{2p-1} < q_{2p} \\ \#\{q_1, \dots, q_{2p}\} = 2p}} F_k(q_1, q_2, \dots, q_{2p-1}, q_{2p}). \quad (7)$$

For $k \geq p$ and $2p < n$,

$$F_{k,p} = \sum_{\substack{l_0, l_1, \dots, l_p \geq 0 \\ l_0 + \dots + l_p = k-p}} Z_{n-2p}^{l_0} \tau_{n-2p+1, n-2p+2} Z_{n-2p+2}^{l_1} \tau_{n-2p+3, n-2p+4} \dots Z_{n-2}^{l_{p-1}} \tau_{n-1, n} Z_n^{l_p} \in \mathcal{M}_p,$$

if $n = 2p$, let

$$F_{k,p} = \sum_{\substack{l_1, \dots, l_p \geq 0 \\ l_1 + \dots + l_p = k-p}} \tau_{1,2} Z_2^{l_1} \tau_{3,4} \cdots Z_{n-2}^{l_{p-1}} \tau_{n-1,n} Z_n^{l_p}.$$

Using this notations, the above formula (7) implies that

$$P_{\mathcal{M}_p} \Delta_{\mathcal{B}_n}^k = \frac{(-1)^{k-p}}{2^p z^p (n-2p)!} \sum_{\sigma \in \mathfrak{S}_n} \sigma F_{k,p} \sigma^{-1}.$$

Using the fact that for $1 \leq i < j \leq n$, $\tau_{j,j+1}$ commutes with Z_i and that elements $(Z_s)_{1 \leq s \leq n}$ commute with each other, we can reorder the product occuring in the expression of $F_{k,p}$. After reordering we get for $2p < n$,

$$F_{k,p} = \tau_{n-2p,n-2p+1} \cdots \tau_{n-1,n} h_{k-p}^{(p+1)}(Z_{n-2p}, Z_{n-b+2}, \dots, Z_n)$$

and if $n = 2p$,

$$F_{k,p} = \tau_{1,2} \cdots \tau_{n-1,n} h_{k-p}^{(p)}(Z_2, Z_4, \dots, Z_n).$$

□

To complete the proof of the formulas, we shall show the following relations in the symmetric group. For any integer p , let $D(\mathfrak{S}_p) = \{\sigma \times \sigma : \sigma \in \mathfrak{S}_p\}$. For any finite group K , we write $\mathcal{P}_K = \frac{1}{|K|} \sum_{k \in K} k \in \mathbb{C}[K]$.

Lemma 4.5. *i) For any integer $i \geq 1$,*

$$(X_1 + X_2 + \cdots + X_{2i}) \mathcal{P}_{\mathcal{H}_{2i}} = \mathcal{P}_{\mathcal{H}_{2i-2}} i(1 + X_{2i-1}) \mathcal{P}_{\mathcal{H}_{2i}}$$

and

$$\sum_{\substack{1 \leq a < b \leq i \\ \text{or } i < a < b \leq 2i}} (a \ b) \mathcal{P}_{D(\mathfrak{S}_i)} = \mathcal{P}_{D(\mathfrak{S}_{i-1})} i X_i \mathcal{P}_{D(\mathfrak{S}_i)}.$$

ii) For any polynomial $P \in \mathbb{C}[z_1, \dots, z_p]$,

$$\mathcal{P}_{\mathcal{H}_{2p}} P(Z_2(z), \dots, Z_{2p}(z)) \mathcal{P}_{\mathcal{H}_{2p}} = \mathcal{P}_{\mathcal{H}_{2p}} P(1, 2(1 + z^{-1} X_3), \dots, p(1 + z^{-1} X_{2p-1})) \mathcal{P}_{\mathcal{H}_{2p}}$$

and

$$\mathcal{P}_{D(\mathfrak{S}_p)} P(Y_1(z), \dots, Y_p(z)) \mathcal{P}_{D(\mathfrak{S}_p)} = \mathcal{P}_{D(\mathfrak{S}_p)} P(1, 2(1 + z^{-1} X_2), \dots, p(1 + z^{-1} X_p)) \mathcal{P}_{D(\mathfrak{S}_p)}.$$

Proof. The assertion ii) follows from the first one. Let us write the left-hand side of the first formula as $\sum_{1 \leq a < b \leq 2i} (a \ b) \mathcal{P}_{\mathcal{H}_{2i}}$. Observe that the transpositions $(2k-1 \ 2k)$ for $1 \leq k \leq i$ belongs to $\mathcal{H}_{2i}$ and that the other transpositions with support in $\{1, \dots, 2i\}$ are conjugated to $(2i-2 \ 2i-1)$ by an element of $\mathcal{H}_{2i-2}$. Using the fact that $X_1 + \cdots + X_{2i} = \sum_{1 \leq a < b \leq 2i} (a \ b)$ is in the center of $\mathfrak{S}_{2i}$, we get the first equality conjugating by elements of $\mathcal{H}_{2i-2}$ on both sides of the equation. The same proof with $D(\mathfrak{S}_k)$ in place of $\mathcal{H}_{2k}$ yields the second formula. □

Proof of Theorem 3.9 Combining Lemma 4.4 and 4.3 yields that for $p < n$,

$$P_{\mathcal{M}_p} \exp[t \Delta_{\mathcal{B}_n}(z)] = \mathcal{I}_{n,t}^p(z)$$

and for $p < n$ or $p = n$ and $n < m$,

$$P_{\mathcal{M}_p} \exp[t\Delta_{\mathcal{B}_{n,m}}(z)] = \mathcal{I}_{n,m,t}^p(z).$$

Moreover, as $\tau_{[1,2p]}$ (resp. $\tilde{\tau}_{[1,p]}$) is invariant by left or right multiplication by elements of $\mathcal{H}_{2p}$ (resp. $D(\mathfrak{S}_p)$), Lemma 4.5 implies that this formulas still hold in the algebra $\mathcal{B}_n(z)$ as $n = 2p$ (resp. in the algebra $\mathcal{B}_{n,n}(z)$ as $n = p$). $\square$

Acknowledgments. The author is grateful to his PhD advisor Thierry Lévy for his insights and his work on a preliminary version of this paper. Many thanks are due to Franck Gabriel and Guillaume Cébron for stimulating discussions.

References

- [1] B. Collins and S. Matsumoto. On some properties of orthogonal Weingarten functions. *Journal of Mathematical Physics*, 50(11):113516, November 2009.
- [2] B. Collins and P. Śniady. Integration with Respect to the Haar Measure on Unitary, Orthogonal and Symplectic Group. *Communications in Mathematical Physics*, 264:773–795, June 2006.
- [3] Benoît Collins. Moments and cumulants of polynomial random variables on unitary groups, the Itzykson-Zuber integral, and free probability. *Int. Math. Res. Not.*, (17):953–982, 2003.
- [4] Benoît Collins and Michael Stolz. Borel theorems for random matrices from the classical compact symmetric spaces. *Ann. Probab.*, 36(3):876–895, 2008.
- [5] C. de Concini and C. Procesi. A characteristic free approach to invariant theory. *Advances in Math.*, 21(3):330–354, 1976.
- [6] R. Goodman and N. Wallach. *Symmetry, Representations, and Invariants*. Springer, 2009.
- [7] Thierry Lévy. Schur-Weyl duality and the heat kernel measure on the unitary group. *Adv. Math.*, 218(2):537–575, 2008.
- [8] I. G. Macdonald. *Symmetric functions and Hall polynomials*. second edition.
- [9] Andrei Okounkov and Anatoly Vershik. A new approach to representation theory of symmetric groups. *Selecta Math. (N.S.)*, 2(4):581–605, 1996.
- [10] P. Zinn-Justin. Jucys-Murphy Elements and Weingarten Matrices. *Letters in Mathematical Physics*, 91:119–127, February 2010.